





























- Framework Directive. Lakes & Reservoirs: Research and Management 8:41-59.
- Randolph K, Wilson J, Tedesco L, Li L, Pascual DL, Soyeux E, 2008. Hyperspectral remote sensing of cyanobacteria in turbid productive water using optically active pigments, chlorophyll a and phycocyanin. *Remote Sens. Environ.* 112: 4009-4019.
- Ravera O, Vollenweider RA, 1968. *Oscillatoria rubescens* D.C. as an indicator of Lago Maggiore eutrophication. *Schweiz. Z. Hydrol.* 30:374-380.
- Reinart A, Reinhold M, 2008. Mapping surface temperature in large lakes with MODIS data. *Remote Sens. Environ.* 112:603-611.
- Ruggiu D, Morabito G, Panzani P, Puggnetti A, 1998. Trends and relations among basic phytoplankton characteristics in the course of the long-term oligotrophication of Lake Maggiore (Italy). *Hydrobiologia* 369-370:243-257.
- Ruiz-Verdú A, Simis SGH, de Hoyos C, Gons HJ, Peña-Martínez R, 2008. An evaluation of algorithms for the remote sensing of cyanobacterial biomass. *Remote Sens. Environ.* 112:3996-4008.
- Salmaso N, Buzzi F, Garibaldi L, Morabito G, Simona M, 2012. Effects of nutrient availability and temperature on phytoplankton development: a case study from large lakes south of the Alps. *Aquat. Sci.* 74:555-570.
- Salmaso N, Mosello R, 2010. Limnological research in the deep southern subalpine lakes: synthesis, directions and perspectives. *Adv. Oceanogr. Limnol.* 1:29-66.
- Santer R, Schmechtig C, 2000. Adjacency effects on water surfaces: primary scattering approximation and sensitivity study. *Appl. Optics* 39:361-375.
- Sass GZ, Creed IF, Bayley SE, Devito KJ, 2007. Understanding variation in trophic status of lakes on the Boreal Plains: a 5-year retrospective using Landsat TM imagery. *Remote Sens. Environ.* 109:127-141.
- Schindler DW, 2006. Recent advances in the understanding and management of eutrophication. *Limnol. Oceanogr.* 51:356-63.
- Schladow SG, Palmarsson SO, Steissberg TE, Hook SJ, Prata FJ, 2004. An extraordinary upwelling event in a deep thermally stratified lake. *Geophys. Res. Lett.* 31:L15504.
- Simis S, Ruiz-Verdú A, Domínguez Gómez J, Peña-Martínez R, Peters S, Gons HJ, 2007. Influence of phytoplankton pigment composition on remote sensing cyanobacterial biomass. *Remote Sens. Environ.* 106:414-427.
- Sobek S, Algesten G, Bergström A-K, Jansson M, Tranvik LJ, 2003. The catchment and climate regulation of pCO<sub>2</sub> in boreal lakes. *Glob. Change Biol.* 9:630-641.
- Song K, Li L, Li Z, Tedesco L, Hall B, Shi K, 2013. Remote detection of cyanobacteria through phycocyanin for water supply source using three-band model. *Ecol. Inform.* 15:22-33.
- Stadelmann TH, Brezonik PL, Kloiber SM, 2001. Seasonal patterns of chlorophyll-a and Secchi disk transparency in lakes of east-central Minnesota: implications for design of ground- and satellite-based monitoring programs. *Lake Reserv. Manage.* 17:299-314.
- Steissberg TE, Hook SJ, Schladow SG, 2005. Characterizing partial upwellings and surface circulation at Lake Tahoe, California-Nevada, USA with thermal infrared images. *Remote Sens. Environ.* 99:2-15.
- Sterckx S, Knaeps E, Ruddick K, 2011. Detection and correction of adjacency effects in hyperspectral airborne data of coastal and inland waters: the use of the near infrared similarity spectrum. *Int. J. Remote Sens.* 32(21): 6479-6505.
- Strömbeck N, Pierson DC, 2001. The effects of variability in the inherent optical properties on estimations of chlorophyll a by remote sensing in Swedish freshwater. *Sci. Total Environ.* 268: 23-137.
- Sugumaran R, Green N, Clayton M, 2007. Monitoring intra-annual water quality variations using airborne hyperspectral remote sensing data in Iowa lakes. *J. Appl. Remote Sens.* 1:013533.
- Tarrant PE, Amacher JA, Neuer S, 2010. Assessing the potential of Medium-Resolution Imaging Spectrometer (MERIS) and Moderate-Resolution Imaging Spectroradiometer (MODIS) data for monitoring total suspended matter in small and intermediate sized lakes and reservoirs. *Water Resour. Res.* 46:W09532.
- Thiemann S, Schiller H, 2003. Determination of the bulk temperature from NOAA/AVHRR satellite data in a midlatitude lake. *Int. J. Appl. Earth. Obs.* 4:339-349.
- Tyler AN, Svab E, Preston T, Présing M, Kovács WA, 2006. Remote sensing of the water quality of shallow lakes: A mixture modeling approach to quantifying phytoplankton in water characterized by high-suspended sediment. *Int. J. Remote Sens.* 27:1521-1537.
- Vincent RK, Qin X, McKay RML, Miner J, Czajkowski K, Savino J, Bridgeman T, 2004. Phycocyanin detection from LANDSAT TM data for mapping cyanobacterial blooms in Lake Erie. *Remote Sens. Environ.* 89:381-392.
- Visconti A, Manca M, de Bernardi R, 2008. Eutrophication-like response to climate warming: an analysis of Lago Maggiore (N. Italy) zooplankton in contrasting years. *J. Limnol.* 67:87-92.
- Wloczyk C, Richter R, Borg E, Neubert W, 2006. Sea and lake surface temperature retrieval from Landsat thermal data in Northern Germany. *Int. J. Remote Sens.* 27:2489-2502.
- Wolter PT, Johnston CA, Niemi GJ, 2005. Mapping submergent aquatic vegetation in the US Great lakes using QuickBird satellite data. *Int. J. Remote Sens.* 26:5255-5274.
- Woźniak SB, Stramski D, 2004. Modeling the optical properties of mineral particles suspended in seawater and their influence on ocean reflectance and chlorophyll estimation from remote sensing algorithms. *Appl. Optics* 43:3489-3503.
- Zhang Y, Pulliainen JT, Koponen SS, Hallikainen MT, 2003. Water quality retrievals from combined Landsat TM data and ERS-2 SAR data in the Gulf of Finland. *IEEE T. Geosci. Remote* 41:622-629.
- Zhou W, Wang S, Zhou Y, Troy A, 2006. Mapping the concentrations of total suspended matter in lake Taihu, China, using Landsat-5 TM data. *Int. J. Remote Sens.* 27:1177-1191.