

A checklist of the freshwater rotifer fauna of Thailand (Rotifera, Monogononta, Bdelloidea)

Phanee SA-ARDRIT,^{1,2} Pornsilp PHOLPUNTHIN,¹ Hendrik SEGERS^{3*}

¹Department of Biology, Prince of Songkla University, 90112 Hat Yai; ²Princess Maha Chakri Sirindhorn Natural History Museum, Prince of Songkla University, 90112 Hat Yai, Thailand; ³Royal Belgian Institute of Natural Sciences, Vautierstraat 29, 1000 Brussels, Belgium

*Corresponding author: hsegers@naturalsciences.be

ABSTRACT

We provide a checklist of the freshwater Rotifera recorded from Thailand, based on a review of available literature. Approximately 398 species of rotifers are recorded. The rotifer fauna of Thailand has been investigated quite comprehensively and these studies focus on four main parts of the country: the northern, north-eastern, central and southern. The number of rotifers on record is the highest in the north-eastern part (275), followed by the southern part (261), the central part (182) and the northern part (115). The majority of Thai Rotifera belongs to family Lecanidae (24.4%), Lepadellidae (11.3%), Brachionidae (11.1%), Trichocercidae (9.0%) and Flosculariidae (9.0%). The most diverse genus is *Lecane* followed by *Trichocerca*, *Lepadella* and *Brachionus*. Although most Thai rotifers are cosmopolitan, there are a number of Oriental endemics, including some strict Thai or regional endemics. Illoricate rotifers and bdelloids are understudied, while experimental and molecular approaches promise to add most to our knowledge and understanding of the role of rotifers in ecosystem functioning.

Key words: Rotifer, biodiversity, biogeography, endemism, Thailand.

Received: February 2013. **Accepted:** June 2013.

INTRODUCTION

Phylum Rotifera is, together with Cladocera and Copepoda, one of three main groups of freshwater zooplankton. In many cases, it contributes the highest number of species to local zooplankton diversity. As with many taxonomic groups, our perception of its diversity has evolved over the years as a consequence of advances in taxonomic knowledge and the availability of new study techniques. This has resulted in the presence of a number of ambiguous records in the literature. In particular, in Thailand rotifers have been recorded over a time span of over a century, starting by Weber (1907), and quite a few older records refer to taxonomic concepts that are now understood differently than at the time they were published, and/or use names that are no longer considered valid.

The application of molecular tools promises to revolutionise our understanding of species-level diversity of many organisms, and this is not different in Rotifera (e.g. Fontaneto *et al.*, 2009; García-Morales and Elías-Gutiérrez, 2013; Leasi *et al.*, 2013). Considering this, we feel that the time has come to re-evaluate available literature records of Thai Rotifera, to serve as a basis for future work utilising the new techniques that are increasingly becoming available. We therefore propose the first checklist of the Rotifera of Thailand, taking into account the most recent insights in the taxonomy and biogeography of the

animals, aligned with contemporary views on taxonomy, diversity and biogeography of Rotifera (Segers 2007, 2008; Segers and De Smet, 2008).

METHODS

Our checklist contains the valid names of genera and species recorded from Thailand, and includes synonyms listed in the relevant literature starting by Weber (1907) and including Meksuwan *et al.* (2011), the most recent paper known to us. Unless stated otherwise, taxonomy and nomenclature follow Segers (2007). We further list infrasubspecific variants recorded under their nominal species, as appropriate. Occurrence of species is indicated by region as defined by Setapan (1999), who recognises six geographical regions (northern, north-eastern, central, western, eastern and southern) defined by mountain ranges and topography in Thailand. For the biogeographical comparison we refer to regions as defined in Balian *et al.* (2008). Information on the distribution of rotifer species is obtained from Segers (1996, 2007), except for those species-level taxa that were, for any reason, not included in those works [*Collotheca algicola* (Hudson, 1886), *Lepadella monostyla* f. *caudata* (Koste, 1972), *Stephanoceros millsii* (Kellicott, 1885), *Trichocerca inermis* (Linder, 1904) and *Trichocerca tenuidens* (Hauer, 1931)].

RESULTS AND DISCUSSION

The study of freshwater rotifers in Thailand started with the publication by Weber (1907). To date, approximately 398 species-level taxa of rotifer have been recorded (Tab. 1). A number of older records were recently re-evaluated by Segers and Savatnalinton (2010). Other dubious or otherwise problematic records are the following:

- *Brachionus dichotomus* Shephard, 1911 and *B. dichotomus reductus* Koste and Shiel, 1980. Both names can be found in literature dealing with Thai Rotifera; however, the typical *B. dichotomus* has, to our knowledge, not been recorded from the Oriental region and we therefore indicate that the presence of *B. dichotomus* Shephard requires confirmation.
- *Brachionus plicatilis* Müller, 1786: records under this name most likely do not refer to this species as presently understood (Ciros-Perez *et al.*, 2001), but any of the warm-water representatives of this cryptic species group (Suatoni *et al.*, 2006).
- *Keratella tropica* Apstein, 1907: we included under this name the record of *Keratella valga* (Ehrenberg, 1834) by Boonsom (1984), considering that the distinction between the cold-water, acidophilic *K. valga* and the warm-water, euryoecious *K. tropica* has long remained problematic, and that it is unlikely that the former would occur in the habitats studied by Boonsom (1984).
- Koste's (1975) record of the nomen nudum *Lepadella monostyla* f. *caudata* (Koste) is most likely a lapsus regarding the taxon *Lepadella monodactyla caudata* Koste, 1972, at present considered a junior subjective synonym of the nominal taxon.
- The presence in Thailand of *Filinia terminalis* (Plate, 1886) is in need of confirmation, considering that this cold-stenotherm is easily confused with the warm-water *F. novaezealandiae* Shiel and Sanoamuang, 1993 (Segers *et al.* 1996). However, while the two records of *F. terminalis* indeed precede the recognition of *F. novaezealandiae*, the presence of this species in the North of Thailand can not be excluded *a priori*.
- The record of *Habrotrocha recumbens* Bartoš, 1963 by Koste (1975) is not included in the checklist. The name is unavailable in the sense of the International Code of Zoological Nomenclature (ICZN, 1999), as it is based solely on the case inhabited by the animal, and not the animal itself.
- Of the remaining species not included in Segers (2007), *Collotheca algicola* (Hudson, 1886) and *Stephanoceros millsii* (Kellicott, 1885) were recently treated as valid by Meksuwan *et al.* (2011), while *Trichocerca inermis* (Linder, 1904) has recently been recognised as a junior synonym of *T. dixonmullalli* (Jennings, 1903) (Jersabek *et al.*, 2011), and *Trichocerca tenuidens* (Hauer, 1931) remains an insufficiently described species *inquirenda* (Segers, 2003).

The majority of Thai Rotifera belongs to family Lecanidae (1 genus with 97 species, 24.4%), Lepadellidae (4 genera with 45 species, 11.3%), Brachionidae (5 genera with 44 species, 11.1%), Trichocercidae (1 genus with 36 species, 9.0%) and Flosculariidae (9 genera with 36 species, 9.0%). The most diverse genus was *Lecane*, comprising 97 species, followed by *Trichocerca* (36 species), *Lepadella* (31 species) and *Brachionus* (30 species).

Regarding the regional distribution of rotifers in the six geographical regions of Thailand (northern, north-eastern, central, western, eastern and southern) (Setapan, 1999), we note that Thailand has been quite comprehensively investigated, but studies have focused on four main parts: the northern, the north-eastern, the central and the southern. The number of rotifers on record is highest in the north-eastern part (275 species) (Sanoamuang *et al.*, 1995; Sanoamuang and Savatnalinton, 1999, 2001; Segers *et al.*, 2004; Savatnalinton and Segers, 2005), followed by the southern part (261 species) (Pholpunthin, 1997; Segers and Pholpunthin, 1997; Pholpunthin and Chittapun, 1998; Chittapun *et al.*, 1999, 2002, 2003, 2005, 2007; Chittapun and Pholpunthin, 2001; Segers and Chittapun, 2001; Sanoamuang, 2007; Meksuwan *et al.*, 2011), the central part (182 species) (De Ridder, 1971; Koste, 1975; Sanoamuang and Segers, 1997; Jithland and Wongrat, 2006; Teeramaethee *et al.*, 2006; Savatnalinton and Segers, 2008; Chittapun *et al.*, 2009; Chittapun, 2011) and the northern part (115 species) (De Ridder, 1971; Sanoamuang, 1998).

That the number of rotifer species recorded from north-east Thailand is the highest of all regions should not automatically be interpreted as indicating that rotifer diversity is the highest there. Admittedly, this region is the largest of all Thai regions, but it is also the most intensively studied region, as can be judged from the number and of the relevant studies and their being comprehensive (*e.g.* Sanoamuang *et al.*, 1995). The southern region has the second largest rotifer record, which we attribute to the high diversity and abundance of its freshwater habitats, in addition to intensive study. The record from other Thai regions is much lower, mostly because the number of studies is, conversely, lower. Nevertheless, when compared to other countries in Southeast Asia, the inventory confirms that the country is the best documented of all countries in the region (Segers, 2001).

The Thai rotifer fauna is composed largely of widespread, cosmopolitan or tropicopolitan species, but there is a sizeable fraction of Oriental (13 species, 3.3%) or even local, Thai endemics (13 species, 3.3%; Tab. 2). It can be expected that the latter number will decrease as research in countries neighbouring Thailand may reveal the presence of these strict Thai endemics, as is the case with species like *Cephalodella songkhlaensis* Segers and Pholpunthin, 1997 and *Ptygura thalenoensis* Meksuwan *et al.*, 2011.

CONCLUSIONS

Notwithstanding that the Thai rotifer fauna is the richest of all Southeast Asian countries, this by no means implies that the fauna is adequately studied. In addition to knowledge gaps concerning understudied regions in Thailand, some taxonomic groups are insufficiently documented. This is the case for taxa such as Notommatidae and bdelloids [respectively 24 and 4 Thai records compared to 48 and 58 from the Oriental region (Segers, 2008)], two groups which are notoriously difficult to study because of methodological constraints. Furthermore, we lack understanding of the ecology and evolution of rotifers, and their role in ecosystem processes (Wallace *et al.*, 2006). Targeted efforts on understudied regions and taxonomic groups, preferably using modern, molecular tools, will undoubtedly raise the records of rotifer diversity in Thailand, while experimental studies are needed to elucidate their contribution to ecosystem processes and functions.

ACKNOWLEDGMENTS

This publication is based on work presented during a workshop on diversity and ecology of freshwater zooplankton in southeast Asia, Guangzhou, China, 2-7 July 2011. We acknowledge grants from the Royal Golden Jubilee PhD Program (No. 4.B.PS/52/D.2), the Faculty of Science, Department of Biology, Prince of Songkla University, and the Institute of Hydrobiology, Jinan University, Guangzhou (Prof. Dr. H. Boping) for the opportunity to attend the workshop. We are most grateful to the organisers of the FISA meeting, Prof. Dr. L. Sanoamuang and Dr. Maria K. Hołyńska, for supporting the participation of HS in the Conference and the opportunity to submit the present contribution to the Proceedings of the meeting.

Finally, we thank two anonymous reviewers for their valuable and constructive comments on this manuscript.

Tab. 1. Species list of Rotifera in Thailand.

Species	Distribution	References
Monogononta		
Family Asplanchnidae		
1 <i>Asplanchna brightwellii</i> Gosse, 1850	NE, C	1, 3, 4, 13, 17, 20, 21, 22
2 <i>Asplanchna priodonta</i> Gosse, 1850	N, NE, C	3, 4, 9, 13, 17, 20, 21, 22, 24
3 <i>Asplanchna sieboldii</i> (Leydig, 1854)	N, NE, C, S	4, 9, 17, 23, 24, 26, 28
4 <i>Asplanchna tropica</i> Koste & Tobias, 1989	NE, S	4, 5, 7
5 <i>Asplanchnopus hyalinus</i> Harring, 1913	NE	4, 17
6 <i>Asplanchnopus multiceps</i> (Schränk, 1793)	NE	4
7 <i>Harringia rousseleti</i> de Beauchamp, 1912	S	23
Family Atrochidae		
8 <i>Acyclus inquietus</i> Leidy, 1882	S	29
9 <i>Cupelopagis vorax</i> (Leidy, 1857)	C, S	2, 5, 7
Family Brachionidae		
10 <i>Anuraeopsis coelata</i> de Beauchamp, 1932	NE, C, S	4, 5, 7, 17, 18, 20, 21, 22, 23, 24
11 <i>Anuraeopsis fissa</i> Gosse, 1851 Syn.: <i>Anuraeopsis fissa fissa</i> Gosse, 1851; Koste, 1975	N, NE, C, S	1, 2, 3, 4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26
12 <i>Anuraeopsis navicula</i> Rousset, 1911	NE, C, S	4, 5, 7, 10, 21, 23, 24, 26
13 <i>Brachionus angularis</i> Gosse, 1851 Incl. <i>Brachionus angularis</i> Gosse f. <i>typica</i> , f. <i>chelonis</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 9, 10, 13, 17, 18, 20, 21, 22, 23, 24, 25, 26
14 <i>Brachionus angularis bidens</i> Plate, 1886	NE	4
15 <i>Brachionus bennini</i> Leissling, 1924	NE	4
16 <i>Brachionus bidentatus</i> Anderson, 1889	NE, C	4, 13, 17, 21, 24
17 <i>Brachionus budapestinensis</i> Daday, 1885	NE	4, 24
18 <i>Brachionus calyciflorus</i> Pallas, 1766 Syn.: <i>Brachionus calyciflorus calyciflorus</i> Pallas, 1766: Sanoamuang <i>et al.</i> , 1995 Incl. <i>Brachionus calyciflorus</i> f. <i>typica</i> , f. <i>monstruosus</i> , f. <i>amphiceros</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 9, 13, 17, 18, 20, 21, 22, 23, 24, 26

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
19 <i>Brachionus caudatus</i> Barrois and Daday, 1894 Incl. <i>Brachionus caudatus</i> f. <i>aculeatus</i> : Sanoamuang <i>et al.</i> , 1995; Teeramaethee <i>et al.</i> , 2006 <i>Brachionus caudatus</i> f. <i>apsteini</i> : Teeramaethee <i>et al.</i> , 2006 <i>Brachionus caudatus</i> f. <i>personatus</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 9, 17, 20, 21, 22, 24, 26
20 <i>Brachionus dichotomus</i> Shephard, 1911	NE, C, S	5, 7, 20, 21, 23, 24
21 <i>Brachionus dichotomus reductus</i> Koste & Shiel, 1980 Syn.: <i>B. dichotomus</i> Shephard f. <i>reductus</i> Koste & Shiel, 1980: Sanoamuang <i>et al.</i> , 1995; Sanoamuang, 1998; Sanoamuang <i>et al.</i> , 2001; Segers <i>et al.</i> , 2004; Savatentalinton & Segers, 2005; Teeramaethee <i>et al.</i> , 2006; Sanoamuang, 2007	N, NE	4, 9, 13, 17, 18, 22
22 <i>Brachionus diversicornis</i> (Daday, 1883)	N, NE, C	3, 4, 9, 17, 20, 21, 22, 24, 26
23 <i>Brachionus donneri</i> Brehm, 1951	N, NE, C, S	3, 4, 5, 7, 9, 13, 17, 20, 22, 23, 24
24 <i>Brachionus durgae</i> Dhanapathi, 1974	NE	4, 24
25 <i>Brachionus falcatus</i> Zacharias, 1898	N, NE, C, S	1, 3, 4, 5, 7, 9, 10, 13, 17, 18, 20, 21, 22, 23, 24, 26, 28
26 <i>Brachionus forficula</i> Wierzejski, 1891 Incl. <i>Brachionus forficula</i> f. <i>typica</i> , f. <i>reductus</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 9, 13, 17, 18, 20, 21, 22, 23, 24, 25, 26, 28
27 <i>Brachionus kostei</i> Shiel, 1983	N, NE	4, 9, 17
28 <i>Brachionus leydigii</i> Cohn, 1862	-	3
29 <i>Brachionus lyratus</i> Shephard, 1911	NE, S	4, 17, 23, 24
30 <i>Brachionus murphyi</i> Sudzuki, 1989 Syn.: <i>Brachionus niwati</i> Sanoamuang, Segers & Dumont, 1995: Sanoamuang <i>et al.</i> , 1995; Pholpunthin, 1997; Segers & Pholpunthin, 1997	NE, S	4, 5, 7, 17, 23
31 <i>Brachionus nilsoni</i> Ahlstrom, 1940	C	22
32 <i>Brachionus plicatilis</i> Müller, 1786	NE	3, 4
33 <i>Brachionus quadridentatus</i> Hermann, 1783 Syn.: <i>Brachionus quadridentatus quadridentatus</i> Hermann, 1783: Sanoamuang <i>et al.</i> , 1995; Athibai <i>et al.</i> , 2005 <i>Brachionus quadridentatus</i> f. <i>typica</i> : Sanoamuang <i>et al.</i> , 1995 Incl. <i>Brachionus quadridentatus</i> f. <i>brevispinus</i> : Sanoamuang <i>et al.</i> , 1995; Athibai <i>et al.</i> , 2005 <i>Brachionus quadridentatus</i> f. <i>cluniorbicularis</i> : Athibai <i>et al.</i> , 2005	N, NE, C, S	1, 3, 4, 5, 7, 9, 10, 13, 15, 17, 18, 20, 21, 22, 23, 24, 26
34 <i>Brachionus quadridentatus melhemi</i> Barrois & Daday, 1894 Syn.: <i>B. quadridentatus</i> f. <i>melhemi</i> Barrois & Daday, 1894: Athibai <i>et al.</i> , 2005	-	18
35 <i>Brachionus quadridentatus mirabilis</i> Daday, 1897 Syn.: <i>B. quadridentatus</i> f. <i>mirabilis</i> Daday, 1897: Chittapun & Pholpunthin, 2001; Chittapun <i>et al.</i> , 2002; Athibai <i>et al.</i> , 2005	S	12, 18
36 <i>Brachionus rotundiformis</i> Tschugunoff, 1921	C, S	18, 19, 23, 26
37 <i>Brachionus rubens</i> Ehrenberg, 1838	NE, C, S	4, 5, 7, 10, 17, 18, 20, 24, 26
38 <i>Brachionus sericus</i> Rousselet, 1907	-	18
39 <i>Brachionus sessilis</i> Varga, 1951	NE	4
40 <i>Brachionus srisumona</i> Segers, Kothetip & Sanoamuang, 2004	NE	17
41 <i>Brachionus urceolaris</i> Müller, 1773	NE, C, S	3, 4, 14, 18, 19, 23, 26, 28
42 <i>Brachionus variabilis</i> Hempel, 1896	S	10
43 <i>Keratella cochlearis</i> (Gosse, 1851) Incl. <i>Keratella cochlearis</i> f. <i>typica</i> , f. <i>micracantha</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 9, 13, 17, 18, 20, 21, 22, 23, 24, 25
44 <i>Keratella edmondsoni</i> Ahlstrom, 1943	NE	4, 13, 17, 18
45 <i>Keratella javana</i> Hauer, 1973	S	12, 15
46 <i>Keratella lenzi</i> Hauer, 1953	N, NE, C, S	2, 4, 5, 7, 9, 13, 17, 18, 20, 21, 22, 24, 26
47 <i>Keratella mixta</i> (Oparina-Charitonova, 1924)	S	15
48 <i>Keratella procurva</i> (Thorpe, 1891)	NE, C	3, 4, 17, 18, 20, 22
49 <i>Keratella taksinensis</i> Chittapun, Pholpunthin & Segers, 2002	S	15
50 <i>Keratella tecta</i> (Gosse, 1851)	NE, C	4, 17, 18, 20, 21, 22

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
51 <i>Keratella tropica</i> (Apstein, 1907) Syn.: <i>Keratella valga tropica</i> Apstein, 1907: Koste, 1975	N, NE, C, S	1, 2, 3, 4, 5, 7, 9, 10, 13, 17, 18, 20, 21, 22, 23, 24, 25, 26
52 <i>Platyonus patulus</i> (Müller, 1786) Syn.: <i>Brachionus patulus</i> Müller, 1786: Boonsom, 1984 <i>Platyonus patulus</i> (Müller) f. <i>typica</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	3, 4, 5, 7, 9, 10, 12, 13, 15, 17, 18, 20, 21, 22, 23, 24, 26
53 <i>Platyias quadricornis</i> (Ehrenberg, 1832) Syn.: <i>Platyias quadricornis quadricornis</i> (Ehrenberg, 1832): Chittapun <i>et al.</i> , 2002 Incl. <i>Platyias quadricornis</i> f. <i>brevispinus</i> : Chittapun <i>et al.</i> , 2002	N, NE, C, S	3, 4, 5, 7, 9, 10, 13, 15, 17, 18, 21, 22, 23, 24, 26
Family Conochilidae		
54 <i>Conochilus (Conochiloides) coenobasis</i> (Skorikov, 1914)	NE	4, 13, 20, 24
55 <i>Conochilus (Conochiloides) dossuarius</i> Hudson, 1885	N, NE	4, 9
56 <i>Conochilus (Conochiloides) natans</i> (Seligo, 1900)	NE, S	4, 13, 20, 23, 24
57 <i>Conochilus (Conochilus) hippocrepis</i> (Schrank, 1803)	NE, S	4, 20, 24, 29
58 <i>Conochilus (Conochilus) unicornis</i> Rousselet, 1892	NE	3, 13
Family Collothecidae		
59 <i>Collothea algicola</i> (Hudson, 1886)	S	29
60 <i>Collothea ambigua</i> (Hudson, 1883)	S	29
61 <i>Collothea campanulata</i> (Dobie, 1849) Incl. <i>Collothea campanulata</i> var. <i>longicaudata</i> : Meksuwan <i>et al.</i> , 2011	NE, S	4, 29
62 <i>Collothea edentata</i> (Collins, 1872)	C	2
63 <i>Collothea heptabrachiata</i> (Schoch, 1869)	S	29
64 <i>Collothea ornata</i> (Ehrenberg, 1832)	S	29
65 <i>Collothea stephanochaeta</i> Edmondson, 1936	S	29
66 <i>Collothea tenuilobata</i> (Anderson, 1889)	NE, S	13, 29
67 <i>Collothea trilobata</i> (Collins, 1872) <i>Collothea</i> cf. <i>trilobata</i> (Collins, 1872): Teeramaethee <i>et al.</i> , 2006	S	29
68 <i>Stephanoceros fimbriatus</i> (Goldfusz, 1820)	S	29
69 <i>Stephanoceros millsii</i> (Kellicott, 1885)	S	29
Family Dicanophoridae		
70 <i>Aspelta circinator</i> (Gosse, 1886)	NE	13
71 <i>Dicanophoroides caudatus</i> (Ehrenberg, 1834) Syn.: <i>Dicanophorus caudatus</i> (Ehrenberg, 1834): Sanoamuang and Savatentalinton, 2001	N, NE	4, 9, 13, 17, 20
72 <i>Dicanophoroides claviger</i> (Hauer, 1965) Syn.: <i>Dicanophorus claviger</i> (Hauer): Pholpunthin, 1997	NE, S	4, 5, 7, 13, 17, 24
73 <i>Dicanophorus epicharis</i> Harring & Myers, 1928 <i>Dicanophorus</i> cf. <i>epicharis</i> Harring & Myers, 1928: Sanoamuang <i>et al.</i> , 1995	NE, C, S	2, 10, 12, 13, 17, 20, 23, 26
74 <i>Dicanophorus grandis</i> (Ehrenberg, 1832)	N, NE	4, 9, 13
75 <i>Dicanophorus prionacis</i> Harring & Myers, 1928	S	5, 7
76 <i>Dicanophorus robustus</i> Harring & Myers, 1928	-	3
77 <i>Encentrum longidens</i> Donner, 1943	S	14
78 <i>Encentrum pornsilpi</i> Segers & Chittapun, 2001	S	14, 19, 23
Family Epiphanidae		
79 <i>Cyrtonia tuba</i> (Ehrenberg, 1834)	S	5, 7
80 <i>Epiphanes clavulata</i> (Ehrenberg, 1832)	N, NE	4, 9, 17, 20, 24
81 <i>Epiphanes macroura</i> (Barrois & Daday, 1894)	NE	4
82 <i>Proalides subtilis</i> Rodewald, 1940	NE	4
83 <i>Proalides tentaculatus</i> de Beauchamp, 1907	NE	4
Family Euchlanidae		
84 <i>Beauchampiella eudactylota</i> (Gosse, 1886) Syn.: <i>Manfredium eudactylotum</i> (Gosse, 1886): Sanoamuang <i>et al.</i> , 1995; Sanoamuang, 1998; Chittapun <i>et al.</i> , 2002; Segers <i>et al.</i> , 2004	N, NE, S	4, 9, 15, 17

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
85 <i>Dipleuchlanis propatula</i> (Gosse, 1886) Incl. <i>Dipleuchlanis propatula</i> f. <i>macroductyla</i> : Chittapun <i>et al.</i> , 2002	N, NE, C, S	3, 4, 5, 7, 9, 12, 13, 15, 17, 20, 23, 24, 26
86 <i>Euchlanis dilatata</i> Ehrenberg, 1832	N, NE, C, S	1, 2, 3, 4, 5, 7, 9, 10, 12, 13, 17, 20, 22, 23, 24, 26
87 <i>Euchlanis incisa</i> Carlin, 1939 Incl. <i>Euchlanis incisa</i> f. <i>mucronata</i> : Chittapun <i>et al.</i> , 1999	N, NE, C, S	4, 9, 10, 12, 13, 15, 17, 20, 22, 24
88 <i>Euchlanis lyra</i> Hudson, 1886 Incl. <i>Euchlanis lyra</i> f. <i>myersi</i> : Chittapun <i>et al.</i> , 1999	S	10
89 <i>Euchlanis meneta</i> Myers, 1930	NE, C, S	2, 10, 13
90 <i>Euchlanis triquetra</i> Ehrenberg, 1838	C	2
91 <i>Tripleuchlanis plicata</i> (Levander, 1894)	N, NE, C	1, 3, 4, 9, 13, 17, 21, 22
Family Flosculariidae		
92 <i>Beauchampia crucigera</i> (Dutrochet, 1812) Syn.: <i>Beauchampia crucigera crucigera</i> (Dutrochet, 1812): Koste, 1975	C, S	2, 29
93 <i>Floscularia armata</i> Segers, 1997	S	29
94 <i>Floscularia bifida</i> Segers, 1997	S	29
95 <i>Floscularia conifera</i> (Hudson, 1886)	NE, C, S	2, 12, 13, 17, 19, 23, 29
96 <i>Floscularia decora</i> Edmondson, 1940	C	2
97 <i>Floscularia melicerta</i> (Ehrenberg, 1832)	C	2
98 <i>Floscularia pedunculata</i> (Joliet, 1883)	S	29
99 <i>Floscularia ringens</i> (Linnaeus, 1758)	NE, C, S	2, 13, 29
100 <i>Floscularia wallacei</i> Segers & Shiel, 2008	S	29
101 <i>Lacinararia flosculosa</i> (Müller, 1773)	S	29
102 <i>Lacinaroides coloniensis</i> (Colledge, 1918)	S	29
103 <i>Limnias ceratophylli</i> Schrank, 1803	C, S	2, 29
104 <i>Limnias melicerta</i> Weisse, 1848	C, S	2, 14, 29
105 <i>Octotrocha speciosa</i> Thorpe, 1893	S	29
106 <i>Pentatrocha gigantea</i> Segers & Shiel, 2008	S	29
107 <i>Ptygura agassizi</i> Edmondson, 1948	S	29
108 <i>Ptygura barbata</i> Edmondson, 1939	S	29
109 <i>Ptygura beauchampi</i> Edmondson, 1940	S	29
110 <i>Ptygura brachiata</i> (Hudson, 1886)	C	2
111 <i>Ptygura crystallina</i> (Ehrenberg, 1834)	S	29
112 <i>Ptygura ctenoida</i> Koste & Tobias, 1990	S	29
113 <i>Ptygura elsteri</i> Koste, 1972 Syn.: <i>Ptygura elsteri elsteri</i> Koste, 1972: Koste, 1975	C, S	2, 29
114 <i>Ptygura furcillata</i> (Kellicott, 1889) Syn.: <i>Ptygura furcillata furcillata</i> (Kellicott, 1889): Koste, 1975 Incl.: <i>Ptygura furcillata</i> f. <i>variabilis</i> : Koste, 1975	NE, C, S	2, 13, 29
115 <i>Ptygura kostei</i> José de Paggi, 1996 Syn.: <i>Ptygura elsteri</i> f. <i>thailandis</i> Koste, 1975: Sanoamuang and Savatnalinton, 2001	NE	13
116 <i>Ptygura longicornis</i> (Davis, 1867)	S	29
117 <i>Ptygura melicerta</i> Ehrenberg, 1832	NE, C	2, 13
118 <i>Ptygura mucicola</i> (Kellicott, 1888)	C, S	2, 29
119 <i>Ptygura noodti</i> (Koste, 1972)	S	29
120 <i>Ptygura pedunculata</i> Edmondson, 1939	S	29
121 <i>Ptygura tacita</i> Edmondson, 1940	NE, S	13, 29
122 <i>Ptygura thalenoensis</i> Meksuwan, Pholpunthin & Segers, 2011	S	29
123 <i>Ptygura wilsonii</i> (Anderson & Shephard, 1892)	S	29
124 <i>Sinantherina ariprepres</i> Edmondson, 1939	NE, C	13, 20, 22
125 <i>Sinantherina semibullata</i> (Thorpe, 1893)	N, NE, S	4, 9, 20, 29
126 <i>Sinantherina socialis</i> (Linnaeus, 1758)	C, S	2, 29
127 <i>Sinantherina spinosa</i> (Thorpe, 1893)	NE, C, S	4, 13, 17, 20, 22, 24, 26, 29

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
Family Gastropodidae		
128 <i>Ascomorpha agilis</i> Zacharias, 1893	C	21
129 <i>Ascomorpha ecaudis</i> Perty, 1850	N, NE, C, S	4, 5, 7, 9, 20, 21
130 <i>Ascomorpha ovalis</i> (Bergendal, 1892)	NE, C, S	3, 4, 7, 13, 17, 20, 21, 22, 23, 24
131 <i>Ascomorpha saltans</i> Bartsch, 1870	NE, C, S	3, 4, 5, 7, 13, 17, 20, 21, 22
132 <i>Gastropus hyptopus</i> (Ehrenberg, 1838)	NE	4
Family Hexarthridae		
133 <i>Hexarthra fennica</i> (Levander, 1892)	NE	4
134 <i>Hexarthra intermedia</i> (Wiszniewski, 1929)	N, NE, C, S	4, 5, 7, 9, 13, 17, 20, 21, 22, 24
135 <i>Hexarthra mira</i> (Hudson, 1871)	N, NE, C, S	4, 5, 7, 9, 13, 17, 19, 20, 22, 23, 24
136 <i>Hexarthra oxyuris</i> (Sernov, 1903)	NE	4
Family Ituridae		
137 <i>Itura aurita</i> (Ehrenberg, 1830)	NE	4
138 <i>Itura symmetrica</i> Segers, Mbogo & Dumont, 1994	NE	4
Family Lecanidae		
139 <i>Lecane abanica</i> Segers, 1994	S	12, 23, 27
140 <i>Lecane acanthinula</i> (Hauer, 1938)	NE, S	4, 6, 14, 23, 27
141 <i>Lecane aculeata</i> (Jakubski, 1912)	N, NE, C, S	2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 27, 28
142 <i>Lecane acus</i> (Harring, 1913)	C	1, 3
143 <i>Lecane aeganea</i> Harring, 1914	N, NE, C	3, 4, 9, 21, 27
144 <i>Lecane agilis</i> (Bryce, 1892)	C	25, 27
145 <i>Lecane arcuata</i> (Bryce, 1891)	NE, C, S	1, 2, 3, 4, 10, 12, 25, 27
146 <i>Lecane arcua</i> Harring, 1914 Syn: <i>Lecane strandi</i> Berzins, 1943: Jithlang & Wongrat, 2006	N, NE, C, S	4, 5, 7, 8, 9, 10, 13, 15, 17, 20, 22, 23, 24, 27
147 <i>Lecane armata</i> Thomasson, 1971	-	27
148 <i>Lecane aspasia</i> Myers, 1917	N, NE	4, 9, 17, 27
149 <i>Lecane baimaii</i> Sanoamuang & Savatentalinton, 1999	NE	11, 17, 27
150 <i>Lecane batillifer</i> (Murray, 1913)	NE, C, S	4, 5, 7, 17, 20, 22, 23, 27
151 <i>Lecane bifastigata</i> Hauer, 1938	NE, C	4, 6, 22, 27
152 <i>Lecane bifurca</i> (Bryce, 1892)	NE, S	5, 7, 10, 19, 20, 23, 24, 27
153 <i>Lecane blachei</i> Bërziņš, 1973	N, NE, C, S	4, 6, 8, 9, 13, 17, 20, 27
154 <i>Lecane braumi</i> Koste, 1988	NE, S	12, 13, 15, 27
155 <i>Lecane bulla</i> (Gosse, 1851) Syn.: <i>Lecane bulla bulla</i> (Gosse, 1851): Segers & Savatentalinton, 2010	N, NE, C, S	1, 2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 19, 20, 21, 22, 23, 24, 26, 27, 28
156 <i>Lecane bulla diabolica</i> (Hauer, 1936)	-	27
157 <i>Lecane calcaria</i> Harring & Myers, 1926	NE, S	7, 12, 17, 27
158 <i>Lecane chinesensis</i> Zhuge & Koste 1996	-	27
159 <i>Lecane clara</i> (Bryce, 1892)	NE, S	5, 7, 12, 20, 27
160 <i>Lecane closteroerca</i> (Schmarda, 1859)	N, NE, C, S	1, 2, 3, 4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27
161 <i>Lecane cornuta</i> (Müller, 1786)	C	1, 2, 3
162 <i>Lecane crenata</i> (Harring, 1913)	C	2, 27
163 <i>Lecane crepida</i> Harring, 1914	N, NE, C, S	2, 4, 5, 7, 9, 10, 12, 13, 17, 20, 22, 23, 24, 27
164 <i>Lecane curvicornis</i> (Murray, 1913) Incl. <i>Lecane curvicornis</i> f. <i>typica</i> : Sanoamuang <i>et al.</i> , 1995	N, NE, C, S	1, 3, 4, 5, 7, 8, 9, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27, 28
165 <i>Lecane decipiens</i> (Murray, 1913)	NE, S	10, 11, 12, 13, 17, 20, 24, 27
166 <i>Lecane donneri</i> Chengalath & Mulamoottil, 1974	NE	11, 17, 24, 27
167 <i>Lecane doryssa</i> Harring, 1914 <i>Lecane</i> cf. <i>doryssa</i> Harring, 1914: Sanoamuang and Savatentalinton, 2001	N, NE, S	4, 9, 12, 13, 15, 17, 20, 23, 24, 27

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
168 <i>Lecane elegans</i> Harring, 1914	N, NE, C	2, 9, 13, 17, 20, 24, 26, 27, 28
169 <i>Lecane enowi</i> Segers and Mertens, 1997	S	15, 27
170 <i>Lecane eswari</i> Dhanapathi, 1976	NE	6, 17, 27
171 <i>Lecane flexilis</i> (Gosse, 1886)	NE, C, S	4, 10, 13, 17, 20, 22, 23, 27
172 <i>Lecane furcata</i> (Murray, 1913)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 27
173 <i>Lecane galeata</i> (Bryce, 1892) Syn.: <i>Lecane (Monostyla) pygmaea</i> Daday, 1897; Koste, 1975	C	2
174 <i>Lecane grandis</i> (Murray, 1913)	NE, S	10, 11, 23, 27
175 <i>Lecane haliclysta</i> Harring & Myers, 1926	N, NE, C, S	4, 9, 12, 13, 17, 20, 22, 23, 24, 27
176 <i>Lecane hamata</i> (Stokes, 1896)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 20, 21, 22, 23, 24, 26, 27
177 <i>Lecane hastata</i> (Murray, 1913)	N, NE, C, S	3, 4, 9, 17, 20, 21, 22, 23, 24, 27
178 <i>Lecane hornemanni</i> (Ehrenberg, 1834)	N, NE, C, S	4, 5, 7, 8, 9, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27
179 <i>Lecane inermis</i> (Bryce, 1892)	NE, C, S	2, 4, 5, 7, 10, 12, 15, 19, 20, 23, 26, 27
180 <i>Lecane inopinata</i> Harring & Myers, 1926	N, NE, C, S	3, 4, 9, 10, 12, 13, 17, 20, 21, 22, 24, 27, 28
181 <i>Lecane intrasinuata</i> (Olofsson, 1917)	C	2
182 <i>Lecane isanensis</i> Sanoamuang & Savatentalinton, 2001	NE	13, 27
183 <i>Lecane junki</i> Koste, 1975	C, S	2, 6, 16, 27
184 <i>Lecane kunthuleensis</i> Chittapun, Pholpunthin & Segers, 2003	S	16, 27
185 <i>Lecane lamellata</i> (Daday, 1893)	-	3
186 <i>Lecane lateralis</i> Sharma, 1978	N, NE, C, S	4, 5, 7, 8, 9, 10, 13, 15, 17, 20, 23, 24, 27, 26, 28
187 <i>Lecane latissima</i> Yamamoto, 1955 Syn.: <i>Lecane thailandensis</i> Segers & Sanoamuang, 1994; Sanoamuang et al., 1995; Sanoamuang and Segers, 1997; Pholpunthin & Chittapun, 1998; Sanoamuang & Savatentalinton, 2001; Segers et al., 2004; Savatentalinton & Segers, 2005	NE, S	4, 6, 8, 13, 17, 20, 27
188 <i>Lecane lauterborni</i> Hauer, 1924	-	3, 27
189 <i>Lecane leontina</i> (Turner, 1892)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27
190 <i>Lecane ludwigii</i> (Eckstein, 1883)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 19, 20, 22, 23, 24, 27
191 <i>Lecane luna</i> (Müller, 1776)	N, NE, C, S	1, 3, 4, 5, 7, 8, 9, 10, 13, 17, 20, 21, 22, 23, 24, 26, 27, 28
192 <i>Lecane lunaris</i> (Ehrenberg, 1832)	N, NE, C, S	2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 25, 27
193 <i>Lecane lungae</i> Savatentalinton & Segers, 2005	NE	20, 27
194 <i>Lecane martensi</i> Savatentalinton & Segers, 2008	C	25, 27
195 <i>Lecane micrognatha</i> Segers & Savatentalinton, 2010	-	27
196 <i>Lecane minuta</i> Segers, 1994	S	5, 6, 7, 27
197 <i>Lecane mitis</i> Harring & Myers, 1926	S	12, 27
198 <i>Lecane monostyla</i> (Daday, 1897)	NE, C, S	3, 4, 10, 12, 15, 17, 22, 23, 24, 27
199 <i>Lecane nana</i> (Murray, 1913)	NE, C, S	1, 3, 4, 5, 7, 13, 17, 20, 27, 28
200 <i>Lecane nelsoni</i> Segers, 1994	NE	13, 27
201 <i>Lecane nitida</i> (Murray, 1913) Syn.: <i>Lecane curvicornis</i> (Murray) f. <i>nitida</i> (Murray): Sanoamuang et al., 1995	NE	4, 27
202 <i>Lecane niwati</i> Segers, Kothetip & Sanoamuang, 2004	NE	17, 27
203 <i>Lecane obtusa</i> (Murray, 1913)	N, NE, C, S	4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 19, 20, 22, 23, 24, 27
204 <i>Lecane opias</i> (Harring & Myers, 1926)	NE	20, 27
205 <i>Lecane palinacis</i> Harring & Myers, 1926	C, S	12, 23, 26, 27, 28
206 <i>Lecane papuana</i> (Murray, 1913)	N, NE, C, S	1, 3, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 24, 26, 27, 28

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
207 <i>Lecane paxiana</i> Hauer, 1940	NE, C	20, 25, 27
208 <i>Lecane pertica</i> Harring & Myers, 1926	N, NE, S	5, 7, 9, 12, 13, 15, 17, 27
209 <i>Lecane punctata</i> (Murray, 1913) Syn.: <i>Lecane harringi</i> Ahlstrom, 1934; Boonsom, 1984; Jithlang and Wongrat, 2006	NE, C	3, 11, 20, 21, 22, 27
210 <i>Lecane pusilla</i> Harring, 1914	N, NE, S	4, 9, 12, 13, 17, 20, 27
211 <i>Lecane pyriformis</i> (Daday, 1905)	NE, C, S	2, 4, 10, 12, 13, 14, 15, 17, 20, 22, 23, 24, 26, 27
212 <i>Lecane quadridentata</i> (Ehrenberg, 1830)	N, NE, C, S	1, 2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 17, 22, 23, 24, 26, 27
213 <i>Lecane rhenana</i> Hauer, 1929	N, NE, C, S	4, 5, 7, 8, 9, 13, 17, 20, 22, 23, 24, 26, 27
214 <i>Lecane rhytida</i> Harring & Myers, 1926	N, NE, C, S	5, 7, 9, 10, 12, 14, 17, 22, 23, 27
215 <i>Lecane robertsonae</i> Segers, 1993	NE, S	17, 20, 23, 27
216 <i>Lecane rutneri</i> Hauer, 1938	NE	4, 13, 17, 20, 24, 27
217 <i>Lecane segersi</i> Sanoamuang, 1996	NE, C, S	6, 14, 17, 20, 23, 26, 27
218 <i>Lecane serrata</i> (Hauer, 1938)	NE	11, 17, 27
219 <i>Lecane shieli</i> Segers & Sanoamuang, 1994	NE, S	4, 6, 10, 13, 17, 20, 27
220 <i>Lecane signifera</i> (Jennings, 1896) f. <i>ploenensis</i> (Voigt, 1902)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27
221 <i>Lecane simonneae</i> Segers, 1993	NE, S	12, 13, 15, 17, 27
222 <i>Lecane sola</i> Hauer, 1936	NE	3, 4, 17, 20, 24, 27
223 <i>Lecane stenroosi</i> (Meissner, 1908)	N, NE, C, S	2, 4, 8, 9, 10, 17, 20, 21, 22, 26, 27
224 <i>Lecane stichaea</i> Harring, 1913	C	2, 21, 27
225 <i>Lecane stichoclysta</i> Segers, 1993	NE	20, 27
226 <i>Lecane subtilis</i> Harring & Myers, 1926	NE, S	17, 23, 27
227 <i>Lecane superaculeata</i> Sanoamuang & Segers, 1997	N, NE, C, S	6, 9, 15, 17, 23, 27
228 <i>Lecane sympoda</i> Hauer, 1929	S	5, 7, 27
229 <i>Lecane syngenes</i> (Hauer, 1938)	S	12, 27
230 <i>Lecane tenuiseta</i> Harring, 1914	NE, C, S	2, 4, 5, 7, 8, 10, 12, 13, 15, 17, 19, 20, 21, 22, 23, 24, 27, 28
231 <i>Lecane thalera</i> (Harring & Myers, 1926)	NE, C	4, 22, 27
232 <i>Lecane thienemanni</i> (Hauer, 1938)	NE, C, S	11, 12, 17, 20, 21, 22, 23, 24, 26, 27
233 <i>Lecane undulata</i> Hauer, 1938	N, NE, C, S	4, 5, 7, 9, 12, 13, 17, 20, 23, 24, 27, 28
234 <i>Lecane unguitata</i> (Fadeev, 1926)	N, NE, C, S	4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 19, 20, 22, 23, 24, 26, 27
235 <i>Lecane ungulata</i> (Gosse, 1887)	N, NE, C, S	2, 4, 5, 7, 8, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26, 27
Family Lepadellidae		
236 <i>Colurella adriatica</i> Ehrenberg, 1831	NE, C, S	1, 3, 4, 12, 13, 20, 23, 24, 25
237 <i>Colurella colurus</i> (Ehrenberg, 1830)	N, NE, C, S	2, 4, 9, 10, 12, 15, 23, 26
238 <i>Colurella colurus compressa</i> (Lucks, 1912)	S	14
239 <i>Colurella hindenburgi</i> Steinecke, 1917	C	21
240 <i>Colurella obtusa</i> (Gosse, 1886)	NE, C, S	2, 4, 5, 7, 10, 12, 13, 14, 15, 17, 20, 21, 23, 24
241 <i>Colurella psammophila</i> Segers & Chittapun, 2001	S	14, 23
242 <i>Colurella sanoamuangae</i> Chittapun, Pholpunthin & Segers, 1999	C, S	10, 14, 23, 26
243 <i>Colurella sulcata</i> (Stenroos, 1898)	NE, S	12, 13, 20, 23
244 <i>Colurella tessellata</i> (Glascott, 1893)	S	12, 23
245 <i>Colurella uncinata</i> (Müller, 1773)	N, NE, C, S	5, 7, 9, 12, 13, 15, 17, 20, 22, 23, 24, 26
246 <i>Colurella uncinata bicuspidata</i> (Ehrenberg, 1832) Syn.: <i>Colurella uncinata</i> (Müller) f. <i>bicuspidata</i> (Ehrenberg): Sanoamuang <i>et al.</i> , 1995; Chittapun <i>et al.</i> , 2002	NE	4
247 <i>Lepadella (Lepadella) acuminata</i> (Ehrenberg, 1834)	NE, C, S	1, 3, 4, 10, 13, 17, 20, 21, 23, 24, 26
248 <i>Lepadella (Lepadella) akrobeles</i> Myers, 1934	NE	11, 17

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
249 <i>Lepadella (Lepadella) amphitropis</i> Harring, 1916	NE	20
250 <i>Lepadella (Lepadella) apsidea</i> Harring, 1916	NE, S	4, 5, 7, 10, 13, 23
251 <i>Lepadella (Lepadella) benjamini</i> Harring, 1916	NE	13, 17
252 <i>Lepadella (Lepadella) biloba</i> Hauer, 1958	NE, S	4, 5, 7, 13, 17
253 <i>Lepadella (Lepadella) costatoides</i> Segers, 1992	N, NE, C, S	4, 9, 13, 17, 20, 22, 23, 24
254 <i>Lepadella (Lepadella) cristata</i> (Rousselet, 1893)	NE, S	12, 13
255 <i>Lepadella (Lepadella) dactyliseta</i> (Stenroos, 1898)	NE, S	4, 5, 7, 12, 24
256 <i>Lepadella (Lepadella) desmeti</i> Segers & Chittapun, 2001	S	14, 23
257 <i>Lepadella (Lepadella) discoidea</i> Segers, 1993	N, NE, C, S	4, 9, 10, 12, 13, 15, 17, 22, 24
258 <i>Lepadella (Lepadella) elliptica</i> Wulfert, 1939	NE	3, 20
259 <i>Lepadella (Lepadella) elongata</i> Koste, 1992 <i>Lepadella</i> cf. <i>elongata</i> Koste, 1992: Sanoamuang & Savatentalinton, 2001	NE	11, 17, 20
260 <i>Lepadella (Lepadella) eurysterna</i> Myers, 1942	NE, S	13, 17, 23
261 <i>Lepadella (Lepadella) latusinus</i> (Hilgendorf, 1899)	N, NE, C, S	2, 4, 5, 7, 9, 13, 23
262 <i>Lepadella (Lepadella) lindau</i> Koste, 1981	NE, C, S	5, 7, 13, 17, 20, 28
263 <i>Lepadella (Lepadella) minoruoides</i> Koste & Robertson, 1983	S	5, 7, 23
264 <i>Lepadella (Lepadella) minuta</i> (Weber & Montet, 1918)	C	25
265 <i>Lepadella (Lepadella) ovalis</i> (Müller, 1786)	N, NE, C, S	2, 3, 4, 5, 7, 9, 10, 12, 13, 17, 20, 22, 23, 24, 26
266 <i>Lepadella (Lepadella) patella</i> (Müller, 1773)	N, NE, C, S	2, 3, 4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 22, 23, 24, 26
267 <i>Lepadella (Lepadella) punctata</i> Wulfert, 1939	S	16
268 <i>Lepadella (Lepadella) quadricarinata</i> (Stenroos, 1898)	N, NE, S	4, 5, 7, 9, 10, 12, 13, 17
269 <i>Lepadella (Lepadella) quinquecostata</i> (Lucks, 1912)	N, NE	9, 13, 17
270 <i>Lepadella (Lepadella) rhomboides</i> (Gosse, 1886)	N, NE, C, S	4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26
271 <i>Lepadella (Lepadella) triba</i> Myers, 1934	NE, C, S	2, 5, 7, 10, 13, 17, 20, 23, 24
272 <i>Lepadella (Lepadella) triptera</i> (Ehrenberg, 1832) Incl.: <i>Lepadella triptera</i> f. <i>alata</i> ; Sanoamuang and Savatentalinton, 2001; Segers, 2004	NE, S	4, 13, 17, 20, 23
273 <i>Lepadella (Lepadella) vandenbrandei</i> Gillard, 1952	N, NE, S	4, 5, 7, 9, 12, 13, 15, 17, 20, 23, 24
274 <i>Lepadella (Heterolepadella) apsicora</i> Myers, 1834	NE, S	4, 5, 7, 13, 15, 17, 20, 23, 24
275 <i>Lepadella (Heterolepadella) ehrenbergii</i> (Perty, 1850)	N, NE, C, S	4, 9, 13, 15, 17, 20, 22, 23
276 <i>Lepadella (Heterolepadella) heterostyla</i> (Murray, 1913)	NE, S	5, 7, 17, 20
277 <i>Lepadella (Xenolepadella) monodactyla</i> Berzins, 1960 Syn.: <i>Lepadella monostyla</i> f. <i>caudata</i> (Koste, 1972)	NE, C, S	2, 13, 15, 23
278 <i>Paracoleurella aemula</i> (Myers, 1934)	S	16
279 <i>Squatinella lamellaris</i> (Müller, 1786) Syn.: <i>Squatinella mutica</i> (Ehrenberg): Chittapun & Pholpunthin, 2001; Chittapun <i>et al.</i> , 2007 Syn.: <i>Squatinella lamellaris</i> (Müller) f. <i>mutica</i> (Ehrenberg): Sanoamuang <i>et al.</i> , 1995; Sanoamuang & Savatentalinton, 2001; Segers <i>et al.</i> , 2004	NE, S	4, 5, 7, 13, 17, 23
280 <i>Squatinella leydigii</i> (Zacharias, 1886) Incl.: <i>Squatinella leydigii</i> f. <i>longiseta</i> : Chittapun <i>et al.</i> , 1999; Chittapun <i>et al.</i> , 2002	S	10, 15
Family Lindiidae		
281 <i>Lindia (Lindia) torulosa</i> Dujardin, 1841	S	19
Family Mytilinidae		
282 <i>Lophocharis salpina</i> (Ehrenberg, 1834)	N, NE, C	4, 9, 13, 17, 20, 24, 26
283 <i>Mytilina acanthophora</i> Hauer, 1938	N, NE	4, 9, 13, 17
284 <i>Mytilina bisulcata</i> (Lucks, 1912)	N, NE, C	3, 9, 13, 17, 22, 24, 26
285 <i>Mytilina compressa</i> (Gosse, 1851)	NE, S	4, 5, 7
286 <i>Mytilina crassipes</i> (Lucks, 1912)	C	21
287 <i>Mytilina michelangellii</i> Reid & Turner, 1988 Syn.: <i>Mytilina ventralis</i> f. <i>diversicantha</i> Wulfert, 1965: Jithlang & Wongrat, 2006	C	21

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
288 <i>Mytilina mucronata</i> (Müller, 1773)	C	21
289 <i>Mytilina unguipes</i> (Lucks, 1912)	N, NE, C	4, 9, 13, 17, 22, 26
290 <i>Mytilina ventralis</i> (Ehrenberg, 1830)	N, NE, C, S	3, 4, 9, 10, 12, 13, 17, 20, 21, 22, 23, 24, 26
291 <i>Mytilina ventralis brevispina</i> (Ehrenberg, 1830)	C	21
Family Notommatidae		
292 <i>Cephalodella forficula</i> (Ehrenberg, 1830) Syn.: <i>Cephalodella forficula forficula</i> (Ehrenberg, 1830): Koste, 1975	NE, C, S	2, 4, 13, 20, 23, 24
293 <i>Cephalodella gibba</i> (Ehrenberg, 1830)	NE, C, S	2, 4, 5, 7, 13, 15, 17, 19, 20, 23, 28
294 <i>Cephalodella</i> cf. <i>hyalina</i> Myers, 1924	S	10
295 <i>Cephalodella innesi</i> Myers, 1924	C, S	10, 14, 15, 19, 23, 28
296 <i>Cephalodella megaloccephala</i> (Glascott, 1893)	S	14
297 <i>Cephalodella mucronata</i> Myers, 1924	NE, C, S	12, 13, 22
298 <i>Cephalodella</i> cf. <i>pachyodon</i> Wulfert, 1937	NE	4
299 <i>Cephalodella plicata</i> Myers, 1924	S	14
300 <i>Cephalodella songkhlaensis</i> Segers & Pholpunthin, 1997	NE, S	7, 20
301 <i>Cephalodella tenuior</i> (Gosse, 1886)	NE, S	13, 23
302 <i>Cephalodella tenuiseta</i> (Burn, 1890)	C	28
303 <i>Cephalodella ventripes</i> (Dixon-Nuttall, 1901)	NE	4
304 <i>Eosphora</i> cf. <i>thoides</i> Wulfert, 1935	NE	4
305 <i>Monommata actices</i> Myers, 1930	NE, S	5, 7, 17
306 <i>Monommata dentata</i> Wulfert, 1940	S	15, 23
307 <i>Monommata grandis</i> Tessin, 1890	S	10, 12, 15, 23
308 <i>Monommata longiseta</i> (Müller, 1786)	S	10, 12, 15
309 <i>Monommata maculata</i> Harring & Myers, 1930	S	12
310 <i>Notommata copeus</i> Ehrenberg, 1834	NE, S	4, 5, 7, 13, 23
311 <i>Notommata pachyura</i> (Gosse, 1886) Incl.: <i>Notommata pachyura</i> f. <i>spinosa</i> : Chittapun & Pholpunthin, 2001	N, NE, S	4, 9, 12, 13, 17, 20
312 <i>Notommata pseudocerberus</i> de Beauchamp, 1908	S	5, 7
313 <i>Notommata pygmaea</i> Harring & Myers, 1922	S	10
314 <i>Notommata saccigera</i> Ehrenberg, 1830	S	12, 15, 23
315 <i>Taphrocampa annulosa</i> Gosse, 1851	S	10, 23
Family Scaridiidae		
316 <i>Scaridium bostjani</i> Daems & Dumont, 1974	N, NE, S	4, 5, 7, 9, 23
317 <i>Scaridium elegans</i> Segers & De Meester, 1994	NE, S	11, 12, 13
318 <i>Scaridium grande</i> Segers, 1995	N, NE, S	9, 12, 13, 15
319 <i>Scaridium longicaudum</i> (Müller, 1786)	N, NE, C, S	4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 22, 23, 24
Family Synchaetidae		
320 <i>Ploesoma lenticulare</i> Herrick, 1885	NE, C	13, 22
321 <i>Ploesoma hudsoni</i> (Imhof, 1891)	NE, C	4, 13, 17, 20, 22
322 <i>Polyarthra euryptera</i> Wierzejski, 1891	-	3
323 <i>Polyarthra longiremis</i> Carlin, 1943	NE	4
324 <i>Polyarthra major</i> Burckhardt, 1900	N, NE	4, 9
325 <i>Polyarthra minor</i> Voigt, 1904	S	5, 7, 10, 15
326 <i>Polyarthra remata</i> Skorikov, 1896	S	5, 7
327 <i>Polyarthra vulgaris</i> Carlin, 1943 <i>Polyarthra</i> cf. <i>vulgaris</i> Carlin, 1943: Sanoamuang, 2007	N, NE, C, S	1, 3, 4, 5, 7, 9, 10, 12, 13, 17, 20, 21, 22, 23, 26
328 <i>Synchaeta longipes</i> Gosse, 1887	NE	4
329 <i>Synchaeta pectinata</i> Ehrenberg, 1832	N, NE	4, 9, 17
330 <i>Synchaeta stylata</i> Wierzejski, 1893	NE, C	4, 13, 17, 20, 21

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
Family Tetrasiphonidae		
331 <i>Tetrasiphon hydrocora</i> Ehrenberg, 1840	S	12
Family Testudinellidae		
332 <i>Pompholyx complanata</i> Gosse, 1851	N, NE, C	1, 3, 4, 9, 17, 20, 21, 22, 24, 26
333 <i>Testudinella ahlstromi</i> Hauer, 1956 Syn.: <i>Testudinella incisa ahlstromi</i> (Hauer): Chittapun & Pholpunthin, 2001	N, NE, S	9, 12, 13, 17, 20, 24
334 <i>Testudinella amphora</i> Hauer, 1938	NE, S	12, 13, 17, 23
335 <i>Testudinella brevicaudata</i> Yamamoto, 1951	N, NE, S	4, 5, 7, 9, 13, 17, 24
336 <i>Testudinella emarginula</i> (Stenroos, 1898)	NE, S	4, 17, 23
337 <i>Testudinella greeni</i> Koste, 1981	N, NE	9, 13, 17, 24
338 <i>Testudinella mucronata</i> (Gosse, 1886)	S	12
339 <i>Testudinella parva</i> (Ternetz, 1892) Syn.: <i>Testudinella insinuata</i> Hauer, 1938; Segers <i>et al.</i> , 2004; <i>T. cf. insinuata</i> Hauer, 1938: Sanoamuang and Savaternalinton, 2001	NE, C, S	4, 12, 13, 15, 17, 20, 21, 24
340 <i>Testudinella patina</i> (Hermann, 1783) Syn.: <i>Testudinella patina</i> (Hermann) f. <i>typica</i> : Sanoamuang <i>et al.</i> , 1995 Incl.: <i>Testudinella patina</i> f. <i>intermedia</i> : Sanoamuang <i>et al.</i> , 1995; Jithlang & Wongrat, 2006	N, NE, C, S	3, 4, 5, 7, 9, 10, 12, 13, 15, 17, 20, 21, 22, 23, 24, 26
341 <i>Testudinella tridentata</i> Smirnov, 1931	N, NE, C, S	4, 9, 12, 13, 15, 17, 20, 24, 26
342 <i>Testudinella walkeri</i> Koste & Shiel, 1980	NE, S	5, 7, 13, 17, 20
Family Trichocercidae		
343 <i>Trichocerca abiloi</i> Segers & Sarma, 1993	NE	13, 17
344 <i>Trichocerca bicristata</i> (Gosse, 1887)	N, NE, C	3, 4, 9, 13, 17, 20, 22, 24
345 <i>Trichocerca bidens</i> (Lucks, 1912)	NE, C, S	4, 13, 17, 20, 22, 23, 24
346 <i>Trichocerca braziliensis</i> (Murray, 1913)	N, NE, C, S	4, 9, 10, 12, 13, 15, 17, 20, 22, 23, 24, 26
347 <i>Trichocerca capucina</i> (Wierzejski & Zacharias, 1893)	N, NE, C, S	4, 9, 13, 17, 20, 21, 22, 23, 24
348 <i>Trichocerca chattoni</i> (de Beauchamp, 1907)	NE, C, S	4, 17, 20, 21, 23, 24
349 <i>Trichocerca collaris</i> (Rousselet, 1896)	NE, S	12, 13
350 <i>Trichocerca cylindrica</i> (Imhof, 1891)	NE, C, S	3, 4, 5, 7, 13, 17, 20, 21, 22, 24
351 <i>Trichocerca dixonnuttalli</i> (Jennings, 1903) Syn.: <i>Trichocerca inermis</i> (Linder, 1904): Sanoamuang and Savaternalinton, 2001; Sanoamuang 2007	NE, S	7, 13, 24
352 <i>Trichocerca elongata</i> (Gosse, 1886)	NE, C	4, 13, 17, 21
353 <i>Trichocerca flagellata</i> Hauer, 1937	N, NE, C, S	4, 5, 7, 9, 12, 13, 17, 22, 23, 24
354 <i>Trichocerca cf. gracilis</i> (Tessin, 1890)	S	23
355 <i>Trichocerca hollaerti</i> De Smet, 1990	NE, S	4, 5, 7, 12, 13, 17, 23, 24
356 <i>Trichocerca insignis</i> (Herrick, 1885)	N, NE, S	4, 5, 7, 9, 12, 13, 17, 20, 24
357 <i>Trichocerca insulana</i> (Hauer, 1937) Syn.: <i>Trichocerca montana</i> Hauer, 1956: Sanoamuang & Savaternalinton, 2001	NE, C, S	4, 10, 13, 20, 23, 24, 26, 28
358 <i>Trichocerca longiseta</i> (Schrank, 1802)	NE, C	13, 17, 20, 21, 24
359 <i>Trichocerca mus</i> Hauer, 1938	C, S	22, 23
360 <i>Trichocerca orca</i> (Murray, 1913)	NE	13
361 <i>Trichocerca porcellus</i> (Gosse, 1851)	NE	11, 13, 17
362 <i>Trichocerca pusilla</i> (Jennings, 1903)	N, NE, C, S	1, 3, 4, 9, 10, 13, 17, 19, 20, 21, 22, 23, 24, 25, 26
363 <i>Trichocerca obtusidens</i> (Olofsson, 1918) Syn.: <i>Trichocerca relicta</i> Donner, 1950: Sanoamuang <i>et al.</i> , 1995; Pholpunthin, 1997; Segers & Pholpunthin, 1997	NE, S	4, 5, 7
364 <i>Trichocerca rosea</i> (Stenroos, 1898)	NE	13
365 <i>Trichocerca rousseleti</i> (Voigt, 1902)	NE	4
366 <i>Trichocerca ruttneri</i> Donner, 1953	NE, S	4, 5, 7, 13, 23
367 <i>Trichocerca scipio</i> (Gosse, 1886) Syn.: <i>Trichocerca jenningsi</i> Voigt, 1957: Sanoamuang & Savaternalinton, 2001; Chittapun <i>et al.</i> , 2002	NE, S	13, 15, 17

To be continued on next page.

Tab. 1. Continued from previous page.

Species	Distribution	References
368 <i>Trichocerca siamensis</i> Segers & Pholpunthin, 1997	NE, S	7, 13, 15, 20
369 <i>Trichocerca similis</i> (Wierzejski, 1893)	N, NE, C, S	3, 4, 5, 7, 9, 10, 13, 17, 20, 21, 22, 23, 24
370 <i>Trichocerca similis grandis</i> Hauer, 1965 Syn.: <i>Trichocerca similis</i> (Weirzejski) f. <i>grandis</i> Hauer: Chittapun & Pholpunthin, 2001; Chittapun <i>et al.</i> , 2002	C, S	12, 15, 26
371 <i>Trichocerca simoneae</i> De Smet 1990	NE	13, 17, 24
372 <i>Trichocerca stylata</i> (Gosse, 1851)	NE, C	4, 13, 17, 22, 24
373 <i>Trichocerca tenuidens</i> (Hauer, 1931)	NE	4
374 <i>Trichocerca tenuior</i> (Gosse, 1886)	N, NE, S	3, 9, 13, 14, 17, 19, 20, 23, 24
375 <i>Trichocerca tigris</i> (Müller, 1786)	NE, S	4, 13, 20, 23
376 <i>Trichocerca voluta</i> (Murray, 1913) Syn.: <i>Trichocerca tropis</i> Hauer, 1937: Sanoamuang <i>et al.</i> , 1995; Pholpunthin, 1997; Segers & Pholpunthin, 1997; Chittapun & Pholpunthin, 2001; Sanoamuang & Savatentalinton, 2001; Segers <i>et al.</i> , 2004	NE, S	4, 5, 7, 12, 13, 17
377 <i>Trichocerca vernalis</i> (Hauer, 1936)	NE	13
378 <i>Trichocerca weberi</i> (Jennings, 1903)	NE, S	13, 17, 20, 23
Family Trichotriidae		
379 <i>Macrochaetus collinsii</i> (Gosse, 1867)	N, NE, C, S	2, 4, 9, 10, 12, 13, 15, 17, 20, 22, 23, 24
380 <i>Macrochaetus danneelae</i> Koste & Shiel, 1983	N, NE	9, 13, 17, 24
381 <i>Macrochaetus longipes</i> Myers, 1934	N, NE, C	4, 9, 13, 17, 22, 24
382 <i>Macrochaetus sericus</i> (Thorpe, 1893)	NE, S	4, 5, 7, 13, 17, 20, 24
383 <i>Macrochaetus subquadratus</i> (Perty, 1850)	NE, C, S	12, 13, 17, 22
384 <i>Trichotria tetractis</i> (Ehrenberg, 1830)	N, NE, C, S	4, 5, 7, 9, 12, 13, 15, 17, 20, 21, 22, 24
385 <i>Wolga spinifera</i> (Western, 1894)	NE	11, 17, 24
Family Trochosphaeridae		
386 <i>Filinia brachiata</i> (Rousselet, 1901)	C	3, 21
387 <i>Filinia camasecla</i> Myers, 1938	N, NE, C	4, 9, 13, 17, 20, 21, 22, 24, 26, 28
388 <i>Filinia longiseta</i> (Ehrenberg, 1834) Incl.: <i>Filinia longiseta</i> var. <i>limnetica</i> : Jithlang & Wongrat, 2006	N, NE, C, S	1, 3, 4, 9, 10, 17, 20, 21, 22, 23, 24, 26
389 <i>Filinia novaezealandiae</i> Shiel & Sanoamuang, 1993	NE, C, S	10, 13, 17, 20, 21, 22, 24, 26, 28
390 <i>Filinia opoliensis</i> (Zacharias, 1898) Syn.: <i>Tetramastix opoliensis</i> Zacharias, 1898: De Ridder, 1971	N, NE, C, S	1, 3, 4, 9, 10, 12, 13, 17, 20, 21, 22, 23, 24, 26
391 <i>Filinia pejlery</i> Hutchinson, 1964	N, NE	4, 9
392 <i>Filinia saltator</i> (Gosse, 1886)	N, NE	4, 9, 17
393 <i>Filinia terminalis</i> (Plate, 1886)	NE	3, 4
394 <i>Trochosphaera aequatorialis</i> Semper, 1872	N, NE, C	3, 4, 9, 17, 26
Bdelloidea		
Family Habrotrochidae		
395 <i>Habrotrocha ampulla</i> Murray, 1911	C	2
396 <i>Habrotrocha angusticollis</i> (Murray, 1905)	C	2
Family Philodinia		
397 <i>Dissotrocha aculeata</i> (Ehrenberg, 1832)	C, S	2, 12, 15, 23
398 <i>Dissotrocha macrostyla</i> (Ehrenberg, 1838)	C	2

N, northern; NE, northeastern; C, central; S, southern; -, no detail about region in publication; 1, De Ridder (1971); 2, Koste (1975); 3, Boonsom (1984); 4, Sanoamuang *et al.* (1995); 5, Pholpunthin (1997); 6, Sanoamuang and Segers (1997); 7, Segers and Pholpunthin (1997); 8, Pholpunthin and Chittapun (1998); 9, Sanoamuang (1998); 10, Chittapun *et al.* (1999); 11, Sanoamuang and Savatentalinton (1999); 12, Chittapun and Pholpunthin (2001); 13, Sanoamuang and Savatentalinton (2001); 14, Segers and Chittapun (2001); 15, Chittapun *et al.* (2002); 16, Chittapun *et al.* (2003); 17, Segers *et al.* (2004); 18, Athibai *et al.* (2005); 19, Chittapun *et al.* (2005); 20, Savatentalinton and Segers (2005); 21, Jithland and Wongrat (2006); 22, Teeramaethee *et al.* (2006); 23, Chittapun *et al.* (2007); 24, Sanoamuang (2007); 25, Savatentalinton and Segers (2008); 26, Chittapun *et al.* (2009); 27, Segers and Savatentalinton (2010); 28, Chittapun (2011); 29, Meksuwan *et al.* (2011).

Tab. 2. List of Oriental and Thai taxa.

Oriental taxa	Thai taxa
1 <i>Brachionus donneri</i> Brehm, 1951	1 <i>Brachionus srisumoniae</i> Segers, Kothetip & Sanoamuang, 2004
2 <i>Brachionus murphyi</i> Sudzuki, 1989	2 <i>Colurella psammophila</i> Segers & Chittapun, 2001
3 <i>Cephalodella songkhlaensis</i> Segers & Pholpunthin, 1997*	3 <i>Encentrum pornsilpi</i> Segers & Chittapun, 2001
4 <i>Colurella sanoamuangae</i> Chittapun, Pholpunthin & Segers, 1999	4 <i>Keratella taksinensis</i> Chittapun, Pholpunthin & Segers, 2002
5 <i>Filinia camasecla</i> Myers, 1938	5 <i>Lecane isanensis</i> Sanoamuang & Savatentalinton, 2001
6 <i>Habrotrocha recumbens</i> Bartoš, 1963	6 <i>Lecane junki</i> Koste, 1975
7 <i>Keratella edmondsoni</i> Ahlstrom, 1943	7 <i>Lecane kunthuleensis</i> Chittapun, Pholpunthin & Segers, 2003
8 <i>Lecane acanthinula</i> (Hauer, 1938)	8 <i>Lecane lungae</i> Savatentalinton & Segers, 2005
9 <i>Lecane blachei</i> Berzins, 1973	9 <i>Lecane martensi</i> Savatentalinton & Segers, 2008
10 <i>Lecane bulla diabolica</i> (Hauer, 1936)	10 <i>Lecane micrognatha</i> Segers & Savatentalinton, 2010
11 <i>Lecane minuta</i> Segers, 1994	11 <i>Lecane niwati</i> Segers, Kothetip & Sanoamuang, 2004
12 <i>Octotrocha speciosa</i> Thorpe, 1893	12 <i>Lecane segersi</i> Sanoamuang, 1996
13 <i>Ptygura thalenoensis</i> Meksuwan, Pholpunthin & Segers, 2011	13 <i>Lecane superaculeata</i> Sanoamuang & Segers, 1997* ^o

*Reliably recorded from Cambodia by Sor (2011); ^o recorded, without substantiation, from Northeast India by Sharma and Sharma (2005).

REFERENCES

- Athibai S, Sanoamuang L, Segers H, 2005. Species diversity of the family Brachionidae (Rotifera) in Thailand. *KKU. Res. J.* 5:1-10.
- Balian EV, Segers H, Lévêque C, Martens K, 2008. An introduction to the Freshwater Animal Diversity Assessment (FADA) project. *Hydrobiologia* 595:3-8.
- Boonsom J, 1984. The freshwater zooplankton of Thailand (Rotifera and Crustacea). *Hydrobiologia* 113:223-229.
- Chittapun S, 2011. Fire and recovery of resting egg bank: an experimental study in paddy fields in Pathum Thani province, Thailand. *Hydrobiologia* 662:163-170.
- Chittapun S, Pholpunthin P, 2001. The rotifer fauna of peat-swamps in southern Thailand. *Hydrobiologia* 446/447:226-269.
- Chittapun S, Pholpunthin P, Segers H, 1999. Rotifera from peat-swamps in Phuket Province, Thailand, with the description of a new *Colurella* Bory de St. Vincent. *Int. Rev. Hydrobiol.* 84:587-593.
- Chittapun S, Pholpunthin P, Segers H, 2002. Rotifer diversity in a peat-swamp in southern Thailand (Narathiwat province) with the description of a new species of *Keratella* Bory de St. Vincent. *Ann. Limnol.-Int. J. Limnol.* 38:185-190.
- Chittapun S, Pholpunthin P, Segers H, 2003. Contribution to the knowledge of Thai microfauna diversity: notes on rare peat swamp Rotifera, with the description of a new *Lecane* Nitzsch, 1872. *Hydrobiologia* 501:7-12.
- Chittapun S, Pholpunthin P, Segers H, 2005. Restoration of tropical peat swamp rotifer communities after perturbation: an experimental study of recovery of rotifers from the resting egg bank. *Hydrobiologia* 546:281-289.
- Chittapun S, Pholpunthin P, Segers H, 2007. Diversity of rotifer fauna from five coastal peat swamps on Phuket Island, Southern Thailand. *ScienceAsia* 33:383-387.
- Chittapun S, Pholpunthin P, Sanoamuang L, 2009. Diversity and composition of zooplankton in rice fields during a crop cycle at Pathum Thani Province, Thailand. *Songklanakarinn J. Sci. Technol.* 31:261-267.
- Ciros-Pérez J, Gómez A, Serra M, 2001. On the taxonomy of three sympatric sibling species of the *Brachionus plicatilis* (Rotifera) complex from Spain, with the description of *B. ibericus* n. sp. *J. Plankton Res.* 23:1311-1328.
- De Ridder M, 1971. [Raderdieren uit het verre Oosten]. [Article in Dutch]. *Biol. Jb. Dodonaea* 39:361-391.
- Fontaneto D, Kaya M, Herniou EA, Barraclough TG, 2009. Extreme levels of hidden diversity in microscopic animals (Rotifera) revealed by DNA taxonomy. *Mol. Phylogenet. Evol.* 53:182-189.
- García-Morales AE, Elías-Gutiérrez M, 2013. DNA barcoding of freshwater Rotifera in Mexico: evidence of cryptic speciation in common rotifers. *Mol. Ecol. Resour.* 13:1097-1107.
- ICZN, 1999. International code of zoological nomenclature. The International Commission on Zoological Nomenclature ed., London: 306 pp.
- Jersabek CD, De Smet WH, Fischer C, Fontaneto D, Michaloudi E, Wallace RL, Segers H, 2011. List of available names in zoology, part phylum Rotifera, species-group names established before 1 January 2000. Available from: http://www.ansp.org/~media/Files/ans/research/sys-rotifera/LAN_CandidatePart-Rotifera-2012-03-22.ashx
- Jithland I, Wongrat L, 2006. Composition and distribution of zooplankton in the Pakak Jolasid Reservoir, Lop Buri Province. *KU. Fish. Res. Bull.* 30:1-18.
- Koste W, 1975. [Über den Rotatorienbestand einer Mikro-biozönose in einem tropischen aquatischen Saumbiotop, der *Eichhornia-crassipes*-Zone im Litoral des Bung-Borapet, einem Stausee in Zentralthailand]. [Article in German]. *Gewässer Abwässer* 57/58:43-58.
- Leasi F, Tang CQ, De Smet WH, Fontaneto D, 2013. Cryptic diversity with wide salinity tolerance in the putative euryhaline *Testudinella clypeata* (Rotifera, Monogononta). *Zool. J. Linn. Soc.-Lond.* 168:17-28.
- Meksuwan P, Pholpunthin P, Segers H, 2011. Diversity of sessile rotifers (Gnesiotrocha, Monogononta, Rotifera) in Thale Noi Lake, Thailand. *Zootaxa* 2997:1-18.

- Pholpunthin P, 1997. Freshwater zooplankton (Rotifera, Cladocera and Copepoda) from Thale-Noi, South Thailand. *J. Sci. Soc. Thailand.* 23:23-34.
- Pholpunthin P, Chittapun S, 1998. Freshwater Rotifera of the genus *Lecane* from Songkhla Province, southern Thailand. *Hydrobiologia* 387/388:23-26.
- Sanoamuang L, 1998. Rotifera of some freshwater habitats in the floodplain of the River Nan, northern Thailand. *Hydrobiologia* 387/388:27-33.
- Sanoamuang L, 2007. The rotifer communities of temporary waters in Northeast Thailand. *KKU Res. J.* 12:203-209.
- Sanoamuang L, Savatentalinton S, 1999. New records of rotifers from Nakhon Ratchasima province, northeast Thailand, with a description of *Lecane baimaii* n. sp. *Hydrobiologia* 412:95-101.
- Sanoamuang L, Savatentalinton S, 2001. The rotifer fauna of Lake Kud-Thing, a shallow lake in Nong Khai Province, northeast Thailand. *Hydrobiologia* 446/447:297-304.
- Sanoamuang L, Segers H, 1997. Additions to the *Lecane* Fauna (Rotifera: Monogononta) of Thailand. *Int. Rev. Ges. Hydrobio.* 82:525-530.
- Sanoamuang L, Segers H, Dumont HJ, 1995. Additions to the rotifer fauna of south-east Asia: new and rare species from north-east Thailand. *Hydrobiologia* 313/314:35-45.
- Savatentalinton S, Segers H, 2005. Rotifers from Kalasin Province, Northeast Thailand, with notes on new and rare species. *Zool. Stud.* 44:361-367.
- Savatentalinton S, Segers H, 2008. Rotifer of waterfall mosses from Phu Hin Rong Kla National Park, Thailand, with the description of *Lecane martensi*, new species (Rotifera: Monogononta: Lecanidae). *Raffles B. Zool.* 56:245-249.
- Segers H, 1996. The biogeography of littoral *Lecane* Rotifera. *Hydrobiologia* 323:169-197.
- Segers H, 2001. Zoogeography of the southeast Asian Rotifera. *Hydrobiologia* 446/447:233-246.
- Segers H, 2003. A biogeographical analysis of rotifers of the genus *Trichocerca* Lamarck, 1801 (Trichocercidae, Monogononta, Rotifera), with notes on taxonomy. *Hydrobiologia* 500:103-114.
- Segers H, 2007. Annotated checklist of the rotifers (Phylum Rotifera), with notes on nomenclature, taxonomy and distribution. *Zootaxa* 1564:1-104.
- Segers H, 2008. Global diversity of rotifers (Phylum Rotifera) in freshwater. *Hydrobiologia* 595:49-59.
- Segers H, Chittapun S, 2001. The interstitial Rotifera of a tropical freshwater peat swamp on Phuket Island, Thailand. *Belg. J. Zool.* 131:65-71.
- Segers H, De Smet WH, 2008. Diversity and endemism in Rotifera: a review, and *Keratella* Bory de St Vincent. *Biodivers. Conserv.* 17:303-316.
- Segers H, Pholpunthin P, 1997. New and rare Rotifera from Thale-Noi Lake, Pattalung Province, Thailand, with a note on the taxonomy of *Cephalodella* (Notommatidae). *Ann. Limnol.-Int. J. Limnol.* 33:13-21.
- Segers H, Savatentalinton S, 2010. A critical re-evaluation of the Lecanidae (Rotifera: Monogononta) of Thailand, with description of a new species. *Int. Rev. Hydrobiol.* 95:343-351.
- Segers H, Koste W, Yussuf SM, 1996. Contribution to the knowledge of the monogonont Rotifera of Zanzibar, with a note on *Filinia novaezealandiae* Shiel & Sanoamuang, 1993. *Int. Rev. Ges. Hydrobio.* 81:597-603.
- Segers H, Kotethip W, Sanoamuang L, 2004. Biodiversity of freshwater microfauna in the floodplain of the River Mun, Northeast Thailand: the Rotifera Monogononta. *Hydrobiologia* 515:1-9.
- Setapan Y, 1999. *Geography of Thailand*. 2. Pisit Publ., Bangkok: 457 pp.
- Sharma BK, Sharma S, 2005. Biodiversity of freshwater rotifers (Rotifera, Eurotatoria) from North-Eastern India. *Zoosyst. Evol.* 81:81-88.
- Sor R, 2011. Comparison of the rotifer Fauna between lakes and reservoirs in the upper part of the Cambodian Mekong River Basin. Degree Thesis. Royal University of Phnom Phen ed., Phnom Phen: 75 pp.
- Suatoni E, Vicario S, Rice S, Snell T, Caccone A, 2006. An analysis of species boundaries and biogeographic patterns in a cryptic species complex: the rotifer-*Brachionus plicatilis*. *Mol. Phylogenet. Evol.* 41:86-98.
- Teeramaethee J, Trevanich A, Sanoamuang L, 2006. Biodiversity and abundance of rotifers in Bueng Boraphet, Nakhon Sawan Province, Thailand. *KKU Res. J.* 11:191-202.
- Wallace RL, Snell TW, Ricci C, Nogrady T, 2006. Rotifera, 1: biology, ecology and systematics. In: H.J. Dumont, T. Nogrady, H. Segers (eds.), *Guides to the identification of the microinvertebrates of the continental waters of the world*. Kenobi Productions and Backhuys Publ.
- Weber EF, 1907. [Rotateurs (Voyage du Dr. Walter Voltz)]. [Article in French]. *Zool. Jahrb. Abt. System.* 24:207-224.