

**Change of macrobenthic communities in the 1930s, 1970s and 2015 in the mesotrophic
Lake Nojiri, Central Japan**

**Kimio HIRABAYASHI,^{1*} Makoto ICHIKAWA,² Shunsuke OKADA,²
Masaru YAMAMOTO³**

¹Division of Applied Biology, Institute of Textile Science and Technology, Academic Assembly, Shinshu University, 3-15-1, Tokida, Ueda City, Nagano Prefecture, 386-8567

²Department of Applied Biology, Faculty of Textile Science and Technology, Shinshu University, 3-15-1, Tokida, Ueda City, Nagano Prefecture, 386-8567

³1-6-12, Yoshimisato-machi, Shimonoseki City, Yamaguchi Prefecture, 759-6525, Japan

*Corresponding author: kimio@shinshu-u.ac.jp

Values of some environmental factors in Lake Nojiri on March 10, 2015

	St. 1	St. 2	St. 3	St. 4	St. 5
Water depth (m)	6.2	10.5	14.9	20.2	27.1
Bottom W. T. (°C)	3.0	2.6	2.7	2.6	2.6
Mud T. (°C)	3.0	3.0	2.6	2.8	2.9
Bottom DO (mg/L)	12.6	11.6	11.8	12.1	12.0
Ignition loss (%)	9.6	11.4	11.2	15.0	13.9